

Exercise sheet 1

Curves and Surfaces, MTH201

1. Find a parametrization $\gamma(t)$ for a line segment joining two given points (x_1, y_1) and (x_2, y_2) . Find $\dot{\gamma}(t)$.
2. What does the parametrization trace out $\gamma(t) = (\frac{2t}{1+t^2}, \frac{1-t^2}{1+t^2})$?
3. Show that the parametrization $\gamma(t) := (t^2 - 1, t(t^2 - 1))$ is not injective, i.e. there are two *distinct* real numbers t_1 and t_2 so that $\gamma(t_1) = \gamma(t_2)$. Can you deduce the shape¹ of this curve? Can you express the set of points defined by this curve as the zero set² of some function $f(x, y)$?
4. For $\mathbf{v} : (\alpha, \beta) \rightarrow \mathbf{R}^2$ and $\mathbf{w} : (\alpha, \beta) \rightarrow \mathbf{R}^2$, show that $(\mathbf{v}(t) \cdot \mathbf{w}(t))' = \mathbf{v}'(t) \cdot \mathbf{w}(t) + \mathbf{v}(t) \cdot \mathbf{w}'(t)$.
5. If $\mathbf{n} : (\alpha, \beta) \rightarrow \mathbf{R}^2$ is such that $\|\mathbf{n}(t)\|$ is constant, then prove that $\dot{\mathbf{n}}(t)$ is either 0 or perpendicular to $\mathbf{n}(t)$.
6. if we denote,

$$s_\alpha(t) := \int_{t_\alpha}^t \|\dot{\gamma}(u)\| du$$

$$s_\beta(t) := \int_{t_\beta}^t \|\dot{\gamma}(u)\| du$$

prove that $s_\beta(t) - s_\alpha(t)$ is a constant (**assume that** $t_\alpha < t_\beta$).

7. If $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ is a smooth **and regular** parametrization, then show that $\|\dot{\gamma}(t)\| : (\alpha, \beta) \rightarrow \mathbb{R}$ is smooth.
8. For the parametrization $\gamma : (-\pi/2, \pi/2) \rightarrow \mathbb{R}^2$ given by $\gamma(t) = (5 \cos(t), 5 \sin(t))$,
 - (a) Find the arc-length function $s(t)$ (starting at, say, 0)
 - (b) Find a reparametrization map ϕ so that $\gamma(\phi(t))$ is a unit-speed parametrization.

¹Just a rough drawing showing where the curve intersects the axes and where it self-intersects etc.

²The zero set of a function $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ is $\{(x, y) \mid f(x, y) = 0\}$, i.e. the set of points (x, y) in the plane for which $f(x, y) = 0$